

Electronic Supplementary Material

Antibody Cocktail Exhibits Broad Neutralization Activity against SARS-CoV-2 and SARS-CoV-2 variants

Yuanyuan Qu^{1#} • Xueyan Zhang^{2,7#} • Meiyu Wang^{3,8#} • Lina Sun^{1#} • Yongzhong Jiang⁴ • Cheng Li⁵ • Wei Wu¹ • Zhen Chen² • Qiangling Yin¹ • Xiaolin Jiang⁶ • Yang Liu¹ • Chuan Li¹ • Jiandong Li¹ • Tianlei Ying⁵ • Dexin Li¹ • Faxian Zhan⁴ • Youchun Wang^{3,8} • Wuxiang Guan^{2✉} • Shiwen Wang^{1✉} • Mifang Liang^{1✉}

1. State Key Laboratory for Molecular Virology and Genetic Engineering, National Institute for Viral Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing 102206, China;
2. Center for Emerging Infectious Diseases, Wuhan Institute of Virology, Chinese Academy of Sciences, Wuhan, Hubei 430071, China;
3. Division of HIV/AIDS and Sex-transmitted Virus Vaccines, Institute for Biological Product Control, National Institutes for Food and Drug Control (NIFDC), Beijing 102629, China;
4. Hubei Provincial Center for Disease Control and Prevention, Wuhan 430065, China;
5. Key Laboratory of Medical Molecular Virology (MOE/NHC/CAMS), School of Basic Medical Sciences, Fudan University, Shanghai 200032, China.
6. Shandong Center for Disease Control and Prevention, Jinan 250014, China.
7. University of Chinese Academy of Sciences, Beijing, 100049, China
8. Peking Union Medical College, Beijing 100730, China

Supporting information to DOI: 10.1007/s12250-021-00409-4




Fig. S1 Characterization of candidates IgGs. **A** The specificity of candidate antibodies detected by FACS. HEK 293T cells expressing SARS-CoV-2 S protein were incubated with candidate antibodies or isotype IgG (HBV mAb) and then stained with anti-human IgG FITC-conjugated antibody. The X-axis represented the fluorescence intensity of human antibodies labeled by FITC. **B** The affinity between antibodies and S1 was measured by BIAcore 8000 system.